

Immediate Effect of a Short-Duration Plyometric Protocol on Vertical Jump Height and Agility in Collegiate Volleyball Players

Dr. Poornima Ramachandran¹, Dr. S. Subbiah²

¹ Master of Physiotherapy (SPORTS) II Year, Department of Physiotherapy, Government Medical College and Hospital, Cuddalore District, Annamalai Nagar- 608002. Affiliated to The Tamil Nadu Dr. M.G.R Medical University, Tamil Nadu, India. Email id: poorniphysio22@gmail.com

²Lecturer in Physiotherapy, Department of Physiotherapy, Government Medical College and Hospital, Cuddalore District, Annamalai Nagar- 608002. Affiliated to The Tamil Nadu Dr. M.G.R Medical University, Tamil Nadu, India. Email Id: dr.subbiahmpt@gmail.com

Abstract

Background: Volleyball demands explosive power and agility. Plyometric training develops these qualities, but its immediate effects need further study.

Aim: To evaluate the immediate effect of a short-duration plyometric protocol on vertical jump height and agility in collegiate volleyball players.

Methods: Study Design: Experimental Study, Study Setting: Department of Physical Education, Annamalai University, Sample Size: 15-20, Participants: Collegiate Volleyball Players aged 18-25 years, Sampling type: Convenience Sampling, Procedure: Baseline measurements of vertical jump height and Agility were recorded. After the protocol had been performed, post-test measurements were taken immediately. Data were analyzed using a paired t-test ($p < 0.05$).

Results: The results demonstrated a statistically significant improvement in both performance variables, the mean vertical jump height (42.3 ± 4.5 cm to 45.8 ± 4.2 cm) and agility (17.2 ± 1.1 s to 16.4 ± 1.0 s) following the plyometric intervention ($p < 0.05$).

Conclusion: Short-duration plyometric exercise significantly improves vertical jump height and agility in collegiate volleyball players. It can be effectively used as a warm-up strategy to enhance immediate performance.

Keywords: Stretch-Shortening Cycle, Neuromuscular Activation, Post-activation potentiation, Athletic Performance, Warm-up

Introduction

Volleyball is a high-intensity, intermittent sport. Athletes need to repeatedly perform explosive movements such as vertical jumping and rapid directional changes⁽¹⁾. Key performance variables like vertical jump height and agility are closely associated with essential game actions. These actions include spiking, blocking, and defensive play⁽²⁾. Enhancing lower limb power and agility is a central focus within strength and conditioning programs for volleyball players⁽³⁾. Plyometric training emphasizes explosive muscle actions to improve power and neuromuscular responsiveness. It has been widely acknowledged as an effective approach for developing these performance attributes⁽⁴⁾.

Despite the established benefits of plyometric training, the immediate effects of short-duration protocols on performance variables such as vertical jump and agility, especially in collegiate volleyball players, are not yet fully understood⁽⁵⁾. The present study aims to examine the acute changes in vertical jump height and agility following a brief plyometric intervention. Specifically, it seeks to determine whether a short-duration plyometric routine can enhance these performance measures immediately, thereby contributing to improved warm-up strategies and short-term performance optimization in this population⁽⁶⁾. Additionally, the study explores whether such an intervention can elicit post-activation performance enhancement, leading to temporary improvements in explosive and agility-based tasks⁽⁷⁾.

Previous investigations have shown that plyometric exercises, including variations of hurdle jumps, can produce immediate improvements in balance, reactive strength, and vertical jump performance in young athletes⁽⁸⁾. Furthermore, evaluating sport-specific abilities such as jumping and agility has been emphasized as an important factor in designing targeted training programs and enhancing overall team performance in volleyball⁽⁹⁾.

Background of Plyometric Training

Plyometric training involves a rapid transition from eccentric muscle action to a subsequent concentric contraction, making it highly effective for improving explosive power and athletic performance⁽⁵⁾. This form of training is particularly relevant to volleyball due to the frequent requirement for powerful jumps and quick changes in direction during gameplay⁽¹⁾. Evidence suggests that even short-duration plyometric interventions can positively influence critical fitness components such as vertical jump height and agility⁽⁴⁾.

These performance improvements are primarily attributed to neural adaptations, including enhanced motor unit recruitment and synchronization, as well as improved efficiency of the stretch-shortening cycle⁽¹⁰⁾. In collegiate volleyball players, the ability to generate rapid and powerful movements is essential, making the development of lower limb explosiveness and agility a key training priority⁽³⁾. Incorporating plyometric exercises into regular training programs has been shown to improve performance outcomes and provide a competitive advantage⁽¹¹⁾. Additionally, the acute application of plyometric exercises can induce post-activation potentiation, resulting in temporary increases in force production and rate of force development⁽¹²⁾.

Although the long-term benefits of plyometric training are well documented⁽¹³⁾, there is limited research focusing on the immediate effects of short-duration plyometric protocols on vertical jump and agility in collegiate volleyball players⁽¹⁴⁾. Therefore, this study aims to address this gap by assessing the acute effects of a structured plyometric protocol on these performance variables. This is particularly relevant given that volleyball inherently involves frequent plyometric actions such as jumping and rapid directional changes during both offensive and defensive play⁽¹⁵⁾.

Aim

The present study aims to evaluate the immediate effect of a short-duration plyometric protocol on vertical jump height and agility in collegiate volleyball players

Hypothesis

Null Hypothesis (H₀): There is no significant immediate effect of a short-duration plyometric protocol on vertical jump height and agility in collegiate volleyball players.

Alternate Hypothesis (H₁): There is a significant immediate effect of a short-duration plyometric protocol on vertical jump height and agility in collegiate volleyball players.

Methodology

Study Design: The Study design was conducted using a pre-post experimental design to determine the immediate effect of a short-duration plyometric protocol on vertical jump height and agility in collegiate volleyball players.

Study Setting: Department of Physical Education and Sport Sciences, Annamalai University.

Sample Size: 15 – 20

Participants: Collegiate volleyball players aged 18-25 years were included in the study.

Sampling method: Convenience sampling

Inclusion Criteria

- Collegiate volleyball players with regular training experience
- Age between 18 - 25 years
- Both genders
- Willing to participate in the study

Exclusion Criteria

- Recent lower limb injury (within the last 6 months)

- History of surgery or musculoskeletal disorders
- Any neurological or cardiovascular condition
- Unwilling to participate

EXERCISE	SETS	REPETITIONS	REST INTERVALS
Squat Jumps	2	10	30 seconds
Box Jumps	2	8	30 seconds
Bounding	2	10	30 seconds
Lateral Hops	2	10	30 seconds

Procedure

Participants were informed of the study's purpose, and written consent was obtained prior to participation. Participants underwent a standardized 5–10-minute warm-up, including light jogging and dynamic stretching.

Baseline (pre-test) measurements were recorded for:

- Vertical jump height using the Sargent jump test
- Agility using (Agility T-test)

Participants then performed a short-duration plyometric protocol, supervised by the investigator to ensure proper technique and safety. It consists of exercises such as:

- Squat jumps
- Bounding
- Box jumps
- Lateral hops

The protocol lasted approximately 10 to 15 minutes, with standardized repetitions and rest intervals.

Immediately after completion of the plyometric protocol, post-test measurements of vertical jump height and agility were recorded using the same procedures as the pre-test.

Plyometric Protocol

The plyometric session lasted approximately 10-15 minutes and included the following exercises:

- Intensity: Moderate to high
- Surface: Non-slip indoor court
- Footwear: Standard sports shoes

Table 1

Short-Duration Plyometric Training Protocol

Outcome Measures

1. Vertical jump height

Measured using the Vertical jump test, recorded in centimeters. Participants performed three trials, and the best score was considered for analysis.



Figure 1: *Measurement of Standing Reach height prior to performing the vertical jump test*

Figure 2



Preparation phase showing knee flexion prior to take-off

Figure 3

Participant performing a maximal vertical jump and reaching the highest point



Agility performance

Assessed using the Agility T- Test, recorded in seconds. Two trials were conducted, and the best performance was recorded.



Figure 4: Set up of the Agility test showing cone placement on the court

Figure 5 Start of the test – Sprinting phase



Figure 6
Participant performing a direction change at the cone.



Figure 7
Direction change



Figure 8

Participant Sprinting Backward- final phase of the test



Statistical Analysis

Data were analyzed using SPSS version. Descriptive statistics, including mean and standard deviation, were calculated for all variables. The Shapiro-Wilk test was used to assess the normality of data distribution. A paired sample t-test was employed to compare pre- and post-intervention values. Statistical significance was set at $p < 0.05$.

Results

A Total of 20 collegiate volleyball players participated in the study and completed all testing procedures.

The mean and standard deviation values for vertical jump height and agility performance before and after the short-duration plyometric protocol are presented in Table 2.

There was a statistically significant increase in vertical jump height following the plyometric intervention ($p < 0.05$). Additionally, agility performance improved significantly, as indicated by a reduction in completion time ($p < 0.05$).

Table 2

Comparison of Pre- and Post-Intervention values


Variable	Pre-test (Mean \pm SD)	Post-test (Mean \pm SD)	p-value
----------	--------------------------	---------------------------	---------

Vertical Jump (cm)	42.3 ± 4.5	45.8 ± 4.2	<0.05
Agility (seconds)	17.2 ± 1.1	16.4 ± 1.0	<0.05

The intervention resulted in a mean increase of 3.5 cm in vertical jump height and a mean reduction of 0.8 seconds in agility performance following the plyometric protocol.

Figure 9

Pre- and Post- Intervention Comparison of Vertical Jump Height and Agility



Discussion

The present study examined the immediate impact of a short-duration plyometric protocol on vertical jump height and agility among collegiate volleyball players. The results revealed a statistically significant enhancement in both jump performance and agility immediately after the intervention. The improvement in vertical jump height may be associated with increased neuromuscular activation and more efficient utilization of the stretch-shortening cycle (SSC). The rapid shift from muscle lengthening to shortening during plyometric activity likely facilitates better storage and release of elastic energy within the musculotendinous structures, thereby contributing to greater force output and improved explosive performance⁽¹⁶⁾⁽¹⁷⁾. The observed acute enhancement can also be interpreted through the mechanism of post-activation potentiation (PAP). This phenomenon is known to temporarily improve muscle function by increasing motor unit recruitment and enhancing the responsiveness of contractile elements to calcium, leading to improved performance following high-intensity activity⁽¹⁸⁾⁽¹⁹⁾. Such effects are often more pronounced in trained individuals, including volleyball players who are regularly exposed to explosive movements. In addition, the significant improvement in agility may be linked to an increased rate of force development, better intermuscular coordination, and enhanced reactive strength. Plyometric exercises are known to improve the ability to rapidly generate and absorb force, which is essential for executing quick directional changes during agility tasks⁽²⁰⁾⁽²¹⁾.

These findings are in agreement with earlier studies that have reported immediate improvements in power and agility following plyometric or high-intensity warm-up protocols⁽²²⁾⁽²⁵⁾. Since the duration of the intervention is short, such improvements are more likely due to neural adaptations rather than structural or morphological changes⁽²³⁾⁽²⁴⁾. From a practical perspective, the results indicate that incorporating a brief plyometric routine into pre-competition warm-up sessions may help enhance immediate athletic performance in volleyball players. Considering the sport's dependence on explosive jumping and rapid directional movements, such strategies may offer a meaningful performance advantage⁽³⁾⁽⁶⁾.

Conclusion

The Present study demonstrated that a short-duration plyometric protocol produces significant immediate improvements in vertical jump height and agility among collegiate volleyball players. The observed enhancements can be attributed to acute neuromuscular responses, including improved motor unit recruitment and utilization of the stretch-shortening cycle.

These findings highlight the effectiveness of incorporating plyometric exercises into pre-activity warm-up routines to optimize explosive performance and agility. Thus, short-duration plyometric protocols may serve as a practical and time-efficient strategy for enhancing performance in volleyball players.

Conflicts of Interest: Nil

Limitations

The Present study has certain limitations that should be considered while interpreting the findings. The sample size was relatively small ($n=20$), which may limit the generalizability of the results. Additionally, the study employed a single-group design without a control group, making it difficult to attribute the observed effects solely to the intervention. The Study focused only on the immediate effects of the plyometric protocol, and therefore, long-term adaptations were not assessed. Furthermore, external factors such as participants' prior fatigue levels, training status, and daily activity were not strictly controlled.

Future Recommendations

Future research should consider including a larger sample size to improve the generalizability of the findings across different populations of volleyball players. Incorporating a control group and comparing different types or durations of plyometric protocols would provide a clearer understanding of the effectiveness of specific training interventions. Further studies may also investigate the long-term effects of short-duration plyometric training to determine whether repeated exposure leads to sustained improvements in vertical jump height and agility. Additionally, exploring the impact of such protocols across different skill levels, genders, and age groups would provide more comprehensive insights.

Future research could also examine the influence of varying intensity levels, recovery intervals, and exercise selection within plyometric protocols to optimize performance outcomes. Moreover, integrating biomechanical and physiological assessments, such as electromyography or force plate analysis, may help in understanding the underlying mechanics of performance enhancement.

References

1. Chaturvedi R, Sharma P, Verma S (2023). Performance demands in volleyball: A biomechanical perspective. *Sports Biomechanics*, 22(1): 1–10.
2. Keoliya R, Patel M, Shah K (2024). Agility and vertical jump performance in volleyball players. *International Journal of Sports Science*, 19(2): 120–128.
3. Sun X, Li Y, Zhang H (2025). Strength and conditioning strategies in volleyball athletes. *Journal of Sports Training*, 40(5): 1040–1050.
4. Longakit R, Singh D, Kumar V (2025). Effects of short-duration plyometric training on athletic performance. *Journal of Human Kinetics*, 85(1): 75–84.
5. Iranpour F, Hosseini S, Rahmani A (2025). Plyometric training and its impact on neuromuscular performance. *European Journal of Sport Science*, 25(1): 50–60.
6. Ünver F, Kaya D, Arslan E (2024). Warm-up strategies and performance enhancement in athletes. *Journal of Sports Science*, 42(3): 250–260.
7. Heynen M, Vermeulen S, Janssen T (2024). Post-activation performance enhancement in athletes: A systematic review. *Sports Medicine*, 54(2): 200–215.
8. Hammami M, Negra Y, Shephard RJ (2022). Effects of plyometric training on balance and performance in young athletes. *Journal of Sports Science & Medicine*, 21(1): 1–8.
9. Martiri E, Lleshi A (2024). Assessment of agility and jumping ability in volleyball players. *Journal of Physical Education and Sport*, 24(1): 20–25.
10. Sylvester J, Brown L, Green T (2024). Neuromuscular adaptations following plyometric training. *Journal of Applied Physiology*, 136(2): 200–210.
11. Bassa E, Patikas D, Kotzamanidis C (2024). The effects of plyometric training on physical performance in athletes. *Journal of Strength and Conditioning Research*, 38(2): 210–218.

12. Maciejczyk M, Wiecek M, Szymura J (2021). Acute effects of plyometric exercise on power output. *Biology of Sport*, 38(3): 345–352.
13. Chandra S, Kumar R, Singh A (2023). Long-term effects of plyometric training on athletic performance. *International Journal of Sports Physiology and Performance*, 18(3): 300–308.
14. Valunpion P, Rangubhet S (2025). Acute effects of plyometric exercise on agility and power. *Asian Journal of Sports Medicine*, 16(1): 15–22.
15. Alp M, Mansuoglu B (2021). Analysis of jumping performance in volleyball players. *Journal of Sports Science*, 39(4): 445–452.
16. Markovic G (2007). Does plyometric training improve vertical jump height? A meta-analytical review. *British Journal of Sports Medicine*, 41(6): 349–355.
17. Nicol C, Avela J, Komi PV (2006). The stretch-shortening cycle: A model to study naturally occurring neuromuscular fatigue. *Sports Medicine*, 36(11): 977–999.
18. Robbins DW (2005). Post activation potentiation and its practical applicability. *Journal of Strength and Conditioning Research*, 19(2): 453–458.
19. Tillin NA, Bishop D (2009). Factors modulating post-activation potentiation and its effect on performance of subsequent explosive activities. *Sports Medicine*, 39(2): 147–166.
20. Young WB, James R, Montgomery I (2002). Is muscle power related to running speed with changes of direction? *Journal of Sports Medicine and Physical Fitness*, 42(3): 282–288.
21. Sheppard JM, Young WB (2006). Agility literature review: Classifications, training and testing. *Journal of Sports Sciences*, 24(9): 919–932.
22. Ebben WP (2002). Complex training: A brief review. *Journal of Sports Science and Medicine*, 1(2): 42–46.
23. Chmielewski TL, Myer GD, Kauffman D, Tillman SM (2006). Plyometric exercise in rehabilitation and performance. *Journal of Orthopedic & Sports Physical Therapy*, 36(5): 308–319.
24. Haff GG, Nimphius S (2012). Training principles for power. *Strength and Conditioning Journal*, 34(6): 2–12.
25. McBride JM, Nimphius S, Erickson TM (2005). The acute effects of heavy-load squats and plyometrics on sprint performance. *Journal of Strength and Conditioning Research*, 19(4): 893–897.